

Bulk and boundary g_2 factorized S -matrices

N. J. MacKay¹ and B. J. Short

*Department of Mathematics,
University of York,
York YO10 5DD, U.K.*

ABSTRACT

We investigate the g_2 -invariant bulk (1+1D, factorized) S -matrix constructed by Ogievetsky, using the bootstrap on the three-point coupling of the vector multiplet to constrain its CDD ambiguity. We then construct the corresponding boundary S -matrix, demonstrating it to be consistent with $Y(g_2, a_1 \times a_1)$ symmetry.

1. Introduction

As a preliminary step in the investigation using tensor methods of 1 + 1-dimensional factorized S -matrices with exceptional $\mathfrak{g}$ (and Yangian $Y(\mathfrak{g})$) invariance, we investigate the case $\mathfrak{g} = g_2$. The factorized S -matrix for the seven-dimensional representation **7** of g_2 was constructed by Ogievetsky [1], and we use this to construct the $g_2 \times g_2$ -invariant S -matrix, applicable in the principal chiral model (PCM). We may choose the S -matrix to have a bootstrap pole for the self-coupling of the **7** multiplet, and the bootstrap applied to this process constrains the CDD factor.

We then investigate the corresponding solutions of the boundary Yang-Baxter equations and the boundary S -matrices for the g_2 PCM — that is, the extension to g_2 of the calculations carried out for classical $\mathfrak{g}$ in [2]. The spectral decomposition is precisely that expected from the $Y(g_2, a_1 \times a_1)$ symmetry [3].

¹email: nm15@york.ac.uk

The method used is the diagrammatic technique of Cvitanovic [4]. We denote the cubic antisymmetric invariant of g_2 as $\succ$, then construct the **7** of g_2 by taking the defining representation of $SO(7)$ and restricting to those $\neg\circ \in SO(7)$ such that $\mathfrak{g}\circ = \succ$. Here $\succ$ satisfies the identities

$$(1.1) \quad \text{---} \diamond \text{---} = -6\text{---}, \quad \begin{array}{c} \diagdown \\ | \\ \diagup \end{array} + \begin{array}{c} \diagup \\ | \\ \diagdown \end{array} = 2 \times - \overline{} -) (\,$$

and it is (sometimes repeated) application of these which is needed to carry out the calculations. If we take the tensor product

$$7 \otimes 7 = 27 \oplus 14 \oplus 7 \oplus 1$$

then the projectors onto the irreducible g_2 -representations are

$$P_{27} = \frac{1}{2} \left(\text{---} + \text{X} \right) - \frac{1}{7} \text{C}, \quad P_{14} = \frac{1}{2} \left(\text{---} - \text{X} \right) + \frac{1}{6} \text{Y},$$

$$P_7 = -\frac{1}{6} \text{Y}, \quad P_1 = \frac{1}{7} \text{C}.$$

2. The bulk S -matrix

The following g_2 -invariant S -matrix satisfies the Yang-Baxter equation [1, 5] :

$$S_{(1,1)}(\theta) = S(\theta) = \sigma(\theta) (P_{\mathbf{27}} + [2]P_{\mathbf{14}} + [8]P_{\mathbf{7}} + [2][12]P_{\mathbf{1}}),$$

where $\sigma(\theta)$ is a scalar prefactor and

$$[y] = \frac{\frac{yi\pi}{12} + \theta}{\frac{yi\pi}{12} - \theta}.$$

Imposing R -matrix unitarity on the S -matrix gives $\sigma(\theta)\sigma(-\theta) = 1$, and imposing hermitian-analyticity gives $\sigma(\theta) = \sigma(-\theta^*)^*$. We rewrite the S -matrix as

$$(2.1) \quad S(\theta) = \omega(\theta) \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} - \frac{6\theta}{i\pi} \text{X} + \frac{2\theta}{(i\pi - \theta)} \right) \left(+ \frac{\theta}{(\frac{2i\pi}{3} - \theta)} \text{Y} \right),$$

where $\sigma(\theta) = (1 - \frac{6\theta}{i\pi})\omega(\theta)$, and impose crossing symmetry,

$$\begin{aligned} \omega(\theta) \left(\text{---} - \frac{6\theta}{i\pi} \text{X} + \frac{2\theta}{(i\pi - \theta)} \right) & \left(+ \frac{\theta}{(\frac{2i\pi}{3} - \theta)} \text{Y} \right) \\ & = \omega(i\pi - \theta) \left(\text{---} \right) \left(- \frac{6(i\pi - \theta)}{i\pi} \text{X} + \frac{2(i\pi - \theta)}{\theta} \text{---} + \frac{(i\pi - \theta)}{(\theta - \frac{i\pi}{3})} \text{Y} \right). \end{aligned}$$

Using $\text{Y} = -\text{Y} + 2\text{X} - \text{---} -$ (we find that this is satisfied if

$$\omega(\theta) = \omega(i\pi - \theta) \frac{(i\pi - \theta)(\frac{2i\pi}{3} - \theta)}{\theta(\frac{i\pi}{3} - \theta)} \Leftrightarrow \sigma(\theta) = \sigma(i\pi - \theta) \frac{(i\pi - \theta)(\frac{2i\pi}{3} - \theta)(\frac{i\pi}{6} - \theta)}{\theta(\frac{i\pi}{3} - \theta)(\theta - \frac{5i\pi}{6})}.$$

To solve for $\sigma(\theta)$ we first introduce

$$\mu_a(\theta) = \frac{\Gamma(\frac{\theta}{2i\pi} + \frac{a}{12}) \Gamma(\frac{-\theta}{2i\pi} + \frac{a}{12} + \frac{1}{2})}{\Gamma(\frac{-\theta}{2i\pi} + \frac{a}{12}) \Gamma(\frac{\theta}{2i\pi} + \frac{a}{12} + \frac{1}{2})},$$

which satisfies $\mu_a(\theta)\mu_a(-\theta) = 1$ and, for real a , $\mu_a(\theta) = \mu_a(-\theta^*)^*$. Further,

$$\frac{\mu_a(\theta)}{\mu_a(i\pi - \theta)} = \frac{(\frac{ai\pi}{6} - \theta)}{(\theta - i\pi + \frac{ai\pi}{6})} = \frac{\mu_{6-a}(i\pi - \theta)}{\mu_{6-a}(\theta)}.$$

We seek a minimal S -matrix, with no poles on the physical strip. The factor $\mu_a(\theta)$ has simple poles at $\theta = -2i\pi n - \frac{ai\pi}{6}$, $\theta = 2i\pi n + i\pi + \frac{ai\pi}{6}$ and simple zeroes at $\theta = 2i\pi n + \frac{ai\pi}{6}$, $\theta = -2i\pi n - i\pi - \frac{ai\pi}{6}$ for $n = 0, 1, 2, \dots$. Thus, to cancel the poles in (2.1), we are led to

$$\sigma(\theta) = \mu_0(-\theta)\mu_1(\theta)\mu_3(\theta)\mu_4(\theta),$$

so that

$$\sigma(\theta) = \frac{\Gamma(\frac{\theta}{2i\pi} + \frac{1}{2}) \Gamma(\frac{-\theta}{2i\pi}) \Gamma(\frac{-\theta}{2i\pi} + \frac{7}{12}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{12}) \Gamma(\frac{-\theta}{2i\pi} + \frac{3}{4}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{4}) \Gamma(\frac{-\theta}{2i\pi} + \frac{5}{6}) \Gamma(\frac{\theta}{2i\pi} + \frac{1}{3})}{\Gamma(\frac{-\theta}{2i\pi} + \frac{1}{2}) \Gamma(\frac{\theta}{2i\pi}) \Gamma(\frac{\theta}{2i\pi} + \frac{7}{12}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{12}) \Gamma(\frac{\theta}{2i\pi} + \frac{3}{4}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{4}) \Gamma(\frac{\theta}{2i\pi} + \frac{5}{6}) \Gamma(\frac{-\theta}{2i\pi} + \frac{1}{3})}$$

(in fact we may choose plus or minus this – our choice of the positive sign will not affect the S -matrix). Thus we have established a minimal S -matrix which is g_2 invariant.

The g_2 PCM S -matrix acts on multiplets which are representations of $g_2 \times g_2$, and is constructed from two minimal S -matrices together with a CDD factor $X(\theta)$:

$$S_{(1,1)}^{PCM}(\theta) = X_{(1,1)}(\theta) \left(S(\theta)_L \otimes S(\theta)_R \right).$$

In order that $S_{(1,1)}^{PCM}(\theta)$ satisfy R -matrix unitarity and crossing-symmetry we require

$$X_{(1,1)}(\theta)X_{(1,1)}(-\theta) = 1 \quad \text{and} \quad \frac{X_{(1,1)}(\theta)}{X_{(1,1)}(i\pi - \theta)} = 1.$$

To construct X we use

$$(y) = (y)_\theta = \frac{\sinh(\frac{\theta}{2} + \frac{yi\pi}{24})}{\sinh(\frac{\theta}{2} - \frac{yi\pi}{24})};$$

this satisfies

$$(y)_\theta(y)_{-\theta} = 1, \quad \frac{(y)_\theta}{(y)_{i\pi - \theta}} = (2y)_{2\theta} \quad \text{and} \quad (y) = (y + 24).$$

The natural choice might be $X = -(2)(4)(8)(10)$, where we have allowed two **7**s to fuse (via simple poles with positive residues) to form either a **7** (at $\theta = 2i\pi/3$) or a $\mathbf{14} \oplus \mathbf{1}$ (at $\theta = i\pi/6$, yielding a multiplet of mass $2 \cos(\pi/12) = \frac{1}{2}(\sqrt{6} + \sqrt{2})$ times the mass of the **7**).

We must then check that the bootstrap equations are satisfied for the scattering of a $\mathbf{7}$ off a fused $\mathbf{7} \subset \mathbf{7} \otimes \mathbf{7}$ (an intricate calculation requiring much repeated application of (1.1)). The minimal S -matrix is consistent with this, but the CDD factor requires an extra factor $(6)^2$, and we must have

$$X_{(1,1)}(\theta) = -(2)(4)(6)^2(8)(10).$$

The apparent double pole at $i\pi/2$ thus introduced is spurious: it is cancelled by a simple zero in each minimal S .

3. The boundary S -matrix

We now consider the half-line case. Following [2], we try a minimal boundary S -matrix of the form

$$K(\theta) = \frac{\tau(\theta)}{(1-c\theta)}(\text{---} + c\theta\text{--}\circ\text{--}).$$

The conditions of boundary R -matrix unitarity and hermitian analyticity impose the constraints

$$(\text{--}\circ\text{--})^\dagger = \text{--}\circ\text{--}, \quad \text{--}\circ\text{--}\circ\text{--} = \text{---}, \quad c \in i\mathbb{R}, \quad \tau(\theta) = \tau(-\theta^*)^* \quad \text{and} \quad \tau(\theta)\tau(-\theta) = 1.$$

We must also impose crossing-unitarity,

$$\begin{aligned} & \frac{\tau(\frac{i\pi}{2}-\theta)}{(1-c(\frac{i\pi}{2}-\theta))} \left(\text{)} + c(\frac{i\pi}{2}-\theta)\text{)} \right) = \\ & \frac{\omega(i\pi-2\theta)\tau(\frac{i\pi}{2}+\theta)}{(1-c(\frac{i\pi}{2}+\theta))} \left(\text{)} \left(-\frac{6(i\pi-2\theta)}{i\pi} \text{X} + \frac{(i\pi-2\theta)}{\theta} \text{---} + \frac{(i\pi-2\theta)}{(2\theta-\frac{i\pi}{3})} \text{X} \right) \text{)} + c(\frac{i\pi}{2}+\theta)\text{)} \right). \end{aligned}$$

After applying (1.1) we find that this implies

$$\begin{aligned} \frac{\tau(\frac{i\pi}{2}-\theta)}{\tau(\frac{i\pi}{2}+\theta)} &= \frac{\omega(i\pi-2\theta)(1-c(\frac{i\pi}{2}-\theta))(\theta-\frac{i\pi}{3})}{(1-c(\frac{i\pi}{2}+\theta))(2\theta-\frac{i\pi}{3})} \left(14 + 2i\pi c\bigcirc + \frac{i\pi}{\theta} + 4 \left(c\bigcirc + \frac{6}{i\pi} \right) \theta \right), \\ \frac{\tau(\frac{i\pi}{2}-\theta)}{\tau(\frac{i\pi}{2}+\theta)} &= \frac{\omega(i\pi-2\theta)(1-c(\frac{i\pi}{2}-\theta))(\theta-\frac{i\pi}{3})(i\pi+2\theta)}{(1-c(\frac{i\pi}{2}+\theta))(2\theta-\frac{i\pi}{3})} \left(\frac{-\alpha}{(\theta-\frac{i\pi}{3})} + \frac{1}{\theta} \mp \frac{12}{i\pi} \right), \end{aligned}$$

together with (for non-trivial $\text{--}\circ\text{--}$) $\text{X} \diamond = \pm \text{)} \text{)}$ and $\text{X} \diamond = \alpha \text{)} \text{)}$ for some constant α . Comparing the two expressions we find $\alpha=0$ and

$$c\bigcirc = -\frac{6(1\pm 1)}{i\pi}.$$

However, $\bigcirc=0$ together with $\text{--}\circ\text{--}\circ\text{--}=\text{---}$ has no solutions in odd dimensions (the eigenvalues of such a matrix are ± 1 , an odd number of which cannot sum to zero). We thus

have $(\text{---}\circ\text{---})^t = \text{---}\circ\text{---}$ and

$$\frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} = \frac{\omega(i\pi - 2\theta)(1 - c(\frac{i\pi}{2} - \theta))(\theta - \frac{i\pi}{3})(i\pi + 2\theta)}{(1 - c(\frac{i\pi}{2} + \theta))(2\theta - \frac{i\pi}{3})} \left(\frac{1}{\theta} - \frac{12}{i\pi} \right),$$

or

$$\frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} = [6] \left[\frac{12}{ci\pi} - 6 \right] \sigma(2\theta), \quad (\text{---}\circ\text{---})^t = \text{---}\circ\text{---} \quad \text{and} \quad c\bigcirc = -\frac{12}{i\pi}.$$

Last we have to impose the boundary Yang-Baxter equation (bYBe). After some algebra we find that this is satisfied if

$$\text{---}\text{---} + \text{---}\text{---}\text{---}\text{---} = \frac{ci\pi}{12} \text{---}\text{---}\text{---}\text{---}.$$

Now using (1.1) we find

$$\text{---}\text{---}\text{---}\text{---} = \text{---}\circ\text{---} + \frac{12}{ci\pi} \text{---}.$$

Thus, putting these two results together,

$$\text{---}\text{---}\text{---}\text{---} = \frac{ci\pi}{12} \text{---}\text{---}\text{---}\text{---} \Leftrightarrow \text{---}\text{---} = \frac{ci\pi}{12} \text{---}\text{---}.$$

Consequently we must have $c = \pm \frac{12}{i\pi}$, with $\text{---}\text{---} = \pm \text{---}\text{---}$ and $\bigcirc = \mp 1$.

In summary, we have shown that the conditions of R -matrix unitarity, hermitian analyticity, crossing unitarity and the bYBe are satisfied by a minimal boundary ‘ K ’-matrix

$$\frac{\tau(\theta)}{(1 \mp \frac{12\theta}{i\pi})} \left(\text{---} \pm \frac{12\theta}{i\pi} \text{---}\text{---} \right) = \tau(\theta) \left(P_- - [\pm 1] P_+ \right), \quad \left(P_{\pm} = \frac{1}{2} (\text{---} \pm \text{---}) \right),$$

where

$$\begin{aligned} (\text{---}\text{---})^\dagger &= \text{---}\text{---}, & (\text{---}\text{---})^t &= \text{---}\text{---}, & \text{---}\text{---}\text{---}\text{---} &= \text{---}, & \text{---}\text{---} &= \pm \text{---}\text{---}, & \bigcirc &= \mp 1, \\ \tau(\theta)\tau(-\theta) &= 1, & \tau(\theta) &= \tau(-\theta^*)^*, & \frac{\tau(\frac{i\pi}{2} - \theta)}{\tau(\frac{i\pi}{2} + \theta)} &= [6] [\pm 1 - 6] \sigma(2\theta). \end{aligned}$$

In fact, since $[1]_{\frac{i\pi}{2} - \theta} / [1]_{\frac{i\pi}{2} + \theta} = [-7] / [-5]$, the choice of sign is redundant – both choices give the same minimal K -matrix. We can write it as

$$\frac{\tau(\theta)}{(1 - \frac{12\theta}{i\pi})} \left(I + \frac{12\theta}{i\pi} E \right) = \tau(\theta) \left(P_- - [1] P_+ \right), \quad \left(P_{\pm} = \frac{1}{2} (I \pm E) \right),$$

where $E = QXQ^{-1}$, $Q \in G_2$, $X = \text{diag}(1, 1, 1, -1, -1, -1, -1)$. This is clearly a subspace of the symmetric space $SO(7)/S(O(3) \times O(4))$; in fact we have

$$E \in \frac{G_2}{SU(2) \times SU(2)},$$

the space of quaternionic subalgebras of the octonions, as may be seen by considering the action of G_2 on a basic triple of octonions [6].

The following constraints are imposed on $\tau(\theta)$:

$$\tau(\theta)\tau(-\theta) = 1, \quad \tau(\theta) = \tau(-\theta^*)^*, \quad \frac{\tau(\frac{i\pi}{2}-\theta)}{\tau(\frac{i\pi}{2}+\theta)} = [6] [-5] \sigma(2\theta).$$

To solve these we note that

$$\frac{\mu_a(\frac{i\pi}{2}-\theta)}{\mu_a(\frac{i\pi}{2}+\theta)} = -[2a-6],$$

and we define

$$\eta_a(\theta) = \frac{\Gamma(\frac{-\theta}{2i\pi} + \frac{a}{12}) \Gamma(\frac{\theta}{2i\pi} + \frac{a}{12} + \frac{1}{4})}{\Gamma(\frac{\theta}{2i\pi} + \frac{a}{12}) \Gamma(\frac{-\theta}{2i\pi} + \frac{a}{12} + \frac{1}{4})}, \quad \text{so that} \quad \frac{\eta_a(\frac{i\pi}{2}-\theta)}{\eta_a(\frac{i\pi}{2}+\theta)} = \mu_{2a-6}(2\theta).$$

This leads us to

$$\tau(\theta) = \mu_{1/2}(\theta)\mu_6(\theta)\eta_{7/2}(\theta)\eta_{9/2}(\theta)\eta_5(\theta)\eta_6(\theta).$$

The simple poles of $\eta_a(\theta)$ are at $\theta = 2i\pi n + \frac{ai\pi}{6}$ and $\theta = -2i\pi n - \frac{i\pi}{2} - \frac{ai\pi}{6}$, while the simple zeroes are at $\theta = -2i\pi n - \frac{ai\pi}{6}$ and $\theta = 2i\pi n + \frac{i\pi}{2} + \frac{ai\pi}{6}$, and so the K -matrix is minimal.

The final piece we require for the complete PCM K -matrix is the factor $Y_1(\theta)$, which must satisfy

$$\frac{Y_1(\frac{i\pi}{2}-\theta)}{Y_1(\frac{i\pi}{2}+\theta)} = X_{(1,1)}(i\pi-2\theta) = X_{(1,1)}(2\theta).$$

We make use of the fact that

$$\frac{(y)_{\frac{i\pi}{2}-\theta}}{(y)_{\frac{i\pi}{2}+\theta}} = (2y)_{i\pi-2\theta} = (2y+24)_{i\pi-2\theta}.$$

Thus the most natural choice is

$$Y_1(\theta) = (1)(2)(-9)^2(-8)(-7)(-6).$$

This has a physical strip simple pole at $\theta = \frac{i\pi}{12}$ at which the minimal K -matrix projects onto the subspace associated with P_+ (the smaller one, and the $(\mathbf{3}, \mathbf{1})$ of $a_1 \times a_1$ as found in [3]). The simple pole at $\theta = \frac{i\pi}{6}$ corresponds to an on-shell diagram which is possible precisely when the bulk three-point coupling of $\mathbf{7}$ s exists.

We should also check the simpler trial solution of [2] for a minimal K -matrix, namely

$$K(\theta) = \rho(\theta) \text{---} \text{---}.$$

Imposing crossing-unitarity gives

$$\frac{\rho(\frac{i\pi}{2}-\theta)}{\rho(\frac{i\pi}{2}+\theta)} \text{---} \text{---} = \omega(i\pi-2\theta) \left(\frac{4(\theta-\frac{i\pi}{3})}{(2\theta-\frac{i\pi}{3})} \text{---} \text{---} + \frac{4(i\pi-2\theta)(i\pi-3\theta)}{i\pi(2\theta-\frac{i\pi}{3})} \text{---} \text{---} + \frac{(i\pi-2\theta)(\theta-\frac{i\pi}{3})}{\theta(2\theta-\frac{i\pi}{3})} \text{---} \text{---} + \frac{(i\pi-2\theta)}{(2\theta-\frac{i\pi}{3})} \text{---} \text{---} \right) \text{---} \text{---},$$

which implies

$$\frac{\rho(\frac{i\pi}{2}-\theta)}{\rho(\frac{i\pi}{2}+\theta)} \rangle = \omega(i\pi-2\theta) \left(\frac{4(\theta-\frac{i\pi}{3})}{(2\theta-\frac{i\pi}{3})} \bigcirc + \frac{4(i\pi-2\theta)(i\pi-3\theta)}{i\pi(2\theta-\frac{i\pi}{3})} \rangle + \frac{(i\pi-2\theta)(\theta-\frac{i\pi}{3})}{\theta(2\theta-\frac{i\pi}{3})} \times + \frac{(i\pi-2\theta)}{(2\theta-\frac{i\pi}{3})} \rangle \right).$$

For non-trivial $\text{---}\bigcirc\text{---}$ we must have $\bigcirc=0$, $(\text{---}\bigcirc\text{---})^t = \pm \text{---}\bigcirc\text{---}$ and $\rangle \times = \alpha \rangle$. But, as pointed out earlier, the constraint $\bigcirc=0$ is inconsistent with $\text{---}\bigcirc\text{---}=\text{---}$. Thus there are no non-trivial solutions of this form.

Acknowledgments: NJM would like to thank Tony Sudbery for a helpful discussion of the octonions, and BJS would like to thank the UK EPSRC for a PhD studentship.

References

- [1] E. Ogievetsky, *Factorized S-matrix with G_2 symmetry*, J.Phys. **G12** (1986) L105
- [2] N. MacKay and B. Short, *Boundary scattering, symmetric spaces and the principal chiral model on the half-line*, Comm. Math. Phys. **233**(2003)313, [hep-th/0104212](#)
- [3] N. MacKay, *Rational K-matrices and representations of twisted Yangians*, J. Phys. **A35** (2002), 7865, [math.QA/0205155](#)
- [4] P. Cvitanovic, *Group theory for Feynman diagrams in non-Abelian gauge theories*, Phys. Rev. **D14** (1976) 1536; *Group Theory: exceptional Lie groups as invariance groups*, web book, <http://www.nbi.dk/GroupTheory/postscript.html>
- [5] E. Ogievetsky and P. Wiegmann, *Factorized S-matrix and the Bethe ansatz for simple Lie groups*, Phys. Lett. **B168** (1986) 360
- [6] J. Baez, *The Octonions*, [math.RA/0105155](#)